

Ramsey Spanning Trees and their Applications


Arnold Filtser

Ben-Gurion University

Co-authors: Ittai Abraham, Shiri Chechik,
Michael Elkin, Ofer Neiman

25 April 2018


Metric Embeddings



Embedding $f : X \rightarrow \mathbb{R}^d$ has distortion α if for all $x, y \in X$

$$d_X(x, y) \leq \|f(x) - f(y)\|_2 \leq \alpha \cdot d_X(x, y)$$

Metric Embeddings




Embedding $f: X \rightarrow \mathbb{R}^d$ has distortion α if for all $x, y \in X$

$$d_X(x, y) \leq \|f(x) - f(y)\|_2 \leq \alpha \cdot d_X(x, y)$$

Theorem (Bourgain,85)

Every n -point metric (X, d_X) is **embeddable** into Euclidean space with **distortion** $O(\log n)$.

Metric Embeddings



Embedding $f: X \rightarrow \mathbb{R}^d$ has distortion α if for all $x, y \in X$

$$d_X(x, y) \leq \|f(x) - f(y)\|_2 \leq \alpha \cdot d_X(x, y)$$


Theorem (Bourgain,85)

Every n -point metric (X, d_X) is **embeddable** into Euclidean space with **distortion** $O(\log n)$.

Asymptotically tight.


Metric Ramsey-Type Problem

For a fixed distortion $k > 1$, what is the largest subset $M \subset X$,
s.t. (M, d_X) is embeddable into Euclidean space with **distortion** k ?



Metric Ramsey-Type Problem


For a fixed distortion $k > 1$, what is the largest subset $M \subset X$,
s.t. (M, d_X) is embeddable into Euclidean space with **distortion** k ?



$$\forall x, y \in M, \quad d_X(x, y) \leq \|f(x) - f(y)\|_2 \leq k \cdot d_X(x, y)$$

Metric Ramsey-Type Problem

For a fixed distortion $k > 1$, what is the largest subset $M \subset X$, s.t. (M, d_X) is embeddable into Euclidean space with **distortion** k ?



$$\forall x, y \in M, \quad d_X(x, y) \leq \|f(x) - f(y)\|_2 \leq k \cdot d_X(x, y)$$

Theorem (Mendel, Naor 07, following BFM86, BLMN05)

For every n -point metric space and $k \geq 1$, there exists a subset M of size $n^{1-1/k}$ that can be embedded into Euclidean space with distortion $O(k)$.

Metric Ramsey-Type Problem

Theorem (Mendel, Naor 07, following BFM86, BLMN05)

For every n -point metric space and $k \geq 1$, there exists a subset M of size $n^{1-1/k}$ that can be embedded into Euclidean space with distortion $O(k)$.

Metric Ramsey-Type Problem

Theorem (Mendel, Naor 07, following BFM86, BLMN05)

For every n -point metric space and $k \geq 1$, there exists a subset M of size $n^{1-1/k}$ that can be embedded into Euclidean space with distortion $O(k)$.

Asymptotically tight.

Metric Ramsey-Type Problem

Theorem (Mendel, Naor 07, following BFM86, BLMN05)

For every n -point metric space and $k \geq 1$, there exists a subset M of size $n^{1-1/k}$ that can be embedded into Euclidean space with distortion $O(k)$.

Asymptotically tight.

Euclidean space can be replaced here by an ultrametric U ! (a.k.a HST)

Metric Ramsey-Type Problem

Theorem (Mendel, Naor 07, following BFM86, BLMN05)

For every n -point metric space and $k \geq 1$, there exists a subset M of size $n^{1-1/k}$ that can be embedded into ultrametric with distortion $O(k)$.

Asymptotically tight.

Euclidean space can be replaced here by an ultrametric U ! (a.k.a HST)

Ultrametric is a special kind of tree which is:

- 1 Very useful for divide and conquer algorithms.
- 2 Isometrically embeds into Euclidean space (i.e. distortion 1).

Our Second Result: Metric Ramsey-Type Problem

Theorem (Mendel, Naor 07, following BFM86, BLMN05)

For every n -point metric space and $k \geq 1$, there exists a subset M of size $n^{1-1/k}$ that can be embedded into ultrametric with distortion $O(k)$.

Our Second Result: Metric Ramsey-Type Problem

Theorem (Mendel, Naor 07, following BFM86, BLMN05)

For every n -point metric space and $k \geq 1$, there exists a subset M of size $n^{1-1/k}$ that can be embedded into ultrametric with distortion $O(128 \cdot k)$.

The constant in the distortion important as it in the exponent.

Our Second Result: Metric Ramsey-Type Problem

Theorem (Mendel, Naor 07, following BFM86, BLMN05)

For every n -point metric space and $k \geq 1$, there exists a subset M of size $n^{1-1/k}$ that can be embedded into ultrametric with distortion $O(128 \cdot k)$.

The constant in the distortion important as it in the exponent.

Paper	Distortion	Size	
BFM06	$O(k \log \log n)$	$n^{1-1/k}$	
BLMN04	$O(k \log k)$	$n^{1-1/k}$	
MN07	$128 \cdot k$	$n^{1-1/k}$	
BGS16	$33 \cdot k$	$n^{1-1/k}$	
NT12	$2e \cdot k$	$n^{1-1/k}$	

Our Second Result: Metric Ramsey-Type Problem

Theorem (Mendel, Naor 07, following BFM86, BLMN05)

For every n -point metric space and $k \geq 1$, there exists a subset M of size $n^{1-1/k}$ that can be embedded into ultrametric with distortion $O(128 \cdot k)$.

The constant in the distortion important as it in the exponent.

Paper	Distortion	Size	
BFM06	$O(k \log \log n)$	$n^{1-1/k}$	Randomized
BLMN04	$O(k \log k)$	$n^{1-1/k}$	Randomized
MN07	$128 \cdot k$	$n^{1-1/k}$	Randomized
BGS16	$33 \cdot k$	$n^{1-1/k}$	Randomized
NT12	$2e \cdot k$	$n^{1-1/k}$	Randomized

Our Second Result: Metric Ramsey-Type Problem

Theorem (Mendel, Naor 07, following BFM86, BLMN05)

For every n -point metric space and $k \geq 1$, there exists a subset M of size $n^{1-1/k}$ that can be embedded into ultrametric with distortion $O(128 \cdot k)$.

The constant in the distortion important as it in the exponent.

Paper	Distortion	Size	
BFM06	$O(k \log \log n)$	$n^{1-1/k}$	Randomized
BLMN04	$O(k \log k)$	$n^{1-1/k}$	Randomized
MN07	$128 \cdot k$	$n^{1-1/k}$	Randomized
BGS16	$33 \cdot k$	$n^{1-1/k}$	Randomized
NT12	$2e \cdot k$	$n^{1-1/k}$	Randomized
This Paper	$8 \cdot k - 2$	$n^{1-1/k}$	Deterministic

Our Second Result: Metric Ramsey-Type Problem

Theorem (Mendel, Naor 07, following BFM86, BLMN05)

For every n -point metric space and $k \geq 1$, there exists a subset M of size $n^{1-1/k}$ that can be embedded into ultrametric with distortion $O(128 \cdot k)$.

The constant in the distortion important as it in the exponent.

Paper	Distortion	Size	
BFM06	$O(k \log \log n)$	$n^{1-1/k}$	Randomized
BLMN04	$O(k \log k)$	$n^{1-1/k}$	Randomized
MN07	$128 \cdot k$	$n^{1-1/k}$	Randomized
BGS16	$33 \cdot k$	$n^{1-1/k}$	Randomized
NT12	$2e \cdot k$	$n^{1-1/k}$	Randomized
This Paper	$8 \cdot k - 2$	$n^{1-1/k}$	Deterministic

*Bartal had similar (deterministic) result.

Our Second Result: Metric Ramsey-Type Problem

Theorem (Our Secondary Result)

For every n -point metric space and $k \geq 1$, there is a **deterministic algorithm** that finds a subset M of size $n^{1-1/k}$ that can be embedded into **ultrametric** with distortion $8 \cdot k$.

Our Second Result: Metric Ramsey-Type Problem

Theorem (Our Secondary Result)

For every n -point metric space and $k \geq 1$, there is a **deterministic algorithm** that finds a subset M of size $n^{1-1/k}$ that can be embedded into ultrametric with distortion $8 \cdot k$.

Instead of preserving distance for $M \times M$,
we can preserve distances for $M \times X$.

Our Second Result: Metric Ramsey-Type Problem

Theorem (Our Secondary Result)

For every n -point metric space and $k \geq 1$, there is a **deterministic algorithm** that finds a subset M of size $n^{1-1/k}$ such that the *half metric* can be embedded into **ultrametric** with distortion $16 \cdot k$ w.r.t $M \times X$.

Instead of preserving distance for $M \times M$,
we can preserve distances for $M \times X$.

Our Second Result: Metric Ramsey-Type Problem

Theorem (Our Secondary Result)

For every n -point metric space and $k \geq 1$, there is a **deterministic algorithm** that finds a subset M of size $n^{1-1/k}$ such that the half metric can be embedded into **ultrametric** with distortion $16 \cdot k$ w.r.t $M \times X$.

Our Second Result: Metric Ramsey-Type Problem

Theorem (Our Secondary Result)

For every n -point metric space and $k \geq 1$, there is a **deterministic algorithm** that finds a subset M of size $n^{1-1/k}$ such that the half metric can be embedded into ultrametric with distortion $16 \cdot k$ w.r.t $M \times X$.

Corollary

For every n -point metric space and $k \geq 1$, there is a **set** \mathcal{U} of $k \cdot n^{\frac{1}{k}}$ **ultrametrics** and a mapping $\text{home} : X \rightarrow \mathcal{U}$, such that for every $x, y \in X$,


$$d_{\text{home}(x)}(x, y) \leq (16 \cdot k) \cdot d_X(x, y)$$

Our Second Result: Metric Ramsey-Type Problem

Corollary

For every n -point metric space and $k \geq 1$,
there is a **set \mathcal{U}** of $k \cdot n^{\frac{1}{k}}$ **ultrametrics** and a mapping
home : $X \rightarrow \mathcal{U}$, such that for every $x, y \in U$,

$$d_{\text{home}(x)}(x, y) \leq (16 \cdot k) \cdot d_X(x, y)$$



Distance Oracle

A distance oracle is a succinct data structure that (approximately) answers distance queries.

Distance Oracle

A distance oracle is a succinct data structure that (approximately) answers distance queries.



Distance Oracle


A distance oracle is a succinct data structure that (approximately) answers distance queries.



The properties of interest are size, distortion and query time.

Distance Oracles: State of the Art

DO	Distortion	Size	Query	Deterministic?
TZ05	$2k - 1$	$O(k \cdot n^{1+1/k})$	$O(k)$	no
MN07	$128k$	$O(n^{1+1/k})$	$O(1)$	no
W13	$(2 + \epsilon)k$	$O(k \cdot n^{1+1/k})$	$O(1/\epsilon)$	no
C14	$2k - 1$	$O(k \cdot n^{1+1/k})$	$O(1)$	no
C15	$2k - 1$	$O(n^{1+1/k})$	$O(1)$	no
RTZ05	$2k - 1$	$O(k \cdot n^{1+1/k})$	$O(k)$	yes
W13	$2k - 1$	$O(k \cdot n^{1+1/k})$	$O(\log k)$	yes



Our contribution: Deterministic Distance Oracles


Distance Oracle	Distortion	Size	Query
RTZ05	$2k - 1$	$O(k \cdot n^{1+1/k})$	$O(k)$
W13	$2k - 1$	$O(k \cdot n^{1+1/k})$	$O(\log k)$
This paper	$8(1 + \epsilon)k$	$O(n^{1+1/k})$	$O(1/\epsilon)$
This paper +C14	$2k - 1$	$O(k \cdot n^{1+1/k})$	$O(1)$

Our contribution: Deterministic Distance Oracles

Corollary

For every n -point metric space and $k \geq 1$,
there is a **set \mathcal{U}** of $k \cdot n^{\frac{1}{k}}$ **ultrametrics** and a mapping
home : $X \rightarrow \mathcal{U}$, such that for every $x, y \in U$,

$$d_{\text{home}(x)}(x, y) \leq (16 \cdot k) \cdot d_X(x, y)$$



Our contribution: Deterministic Distance Oracles

Corollary

For every n -point metric space and $k \geq 1$,
there is a **set \mathcal{U}** of $k \cdot n^{\frac{1}{k}}$ **ultrametrics** and a mapping
home : $X \rightarrow \mathcal{U}$, such that for every $x, y \in U$,

$$d_{\text{home}(x)}(x, y) \leq (16 \cdot k) \cdot d_X(x, y)$$



Our contribution: Deterministic Distance Oracles

Corollary

For every n -point metric space and $k \geq 1$,
there is a **set** \mathcal{U} of $k \cdot n^{\frac{1}{k}}$ **ultrametrics** and a mapping
home : $X \rightarrow \mathcal{U}$, such that for every $x, y \in U$,

$$d_{\text{home}(x)}(x, y) \leq (16 \cdot k) \cdot d_X(x, y)$$

Theorem (Tree Distance Oracle, HT84, BFC00)

For every **tree metric**, there is an exact distance oracle of **linear size** and **constant query time**.

Our contribution: Deterministic Distance Oracles

Corollary

For every n -point metric space and $k \geq 1$,
there is a **set** \mathcal{U} of $k \cdot n^{\frac{1}{k}}$ **ultrametrics** and a mapping
home : $X \rightarrow \mathcal{U}$, such that for every $x, y \in U$,

$$d_{\text{home}(x)}(x, y) \leq (16 \cdot k) \cdot d_X(x, y)$$

Theorem (Tree Distance Oracle, HT84, BFC00)

For every **tree metric**, there is an exact distance oracle of **linear size** and **constant query time**.

Theorem (Ramsey based Deterministic Distance Oracle)

For any n -point metric space, there is a distance oracle with :

Distortion	Size	Query time
$16 \cdot k$	$O(k \cdot n^{1+1/k})$	$O(1)$

Our contribution: Deterministic Distance Oracles

Corollary

For every n -point metric space and $k \geq 1$,
there is a **set \mathcal{U}** of $k \cdot n^{\frac{1}{k}}$ **ultrametrics** and a mapping
home : $X \rightarrow \mathcal{U}$, such that for every $x, y \in U$,

$$d_{\text{home}(x)}(x, y) \leq (16 \cdot k) \cdot d_X(x, y)$$

Theorem (Ramsey based Deterministic Distance Oracle)

For any n -point metric space, there is a distance oracle with :

Distance Oracle	Distortion	Size	Query
This paper	$16 \cdot k$	$O(k \cdot n^{1+1/k})$	$O(1)$
This paper	$8(1 + \epsilon)k$	$O(n^{1+1/k})$	$O(1/\epsilon)$
This paper + C14	$2k - 1$	$O(k \cdot n^{1+1/k})$	$O(1)$

Our contribution: Deterministic Distance Oracles

Corollary

For every n -point metric space and $k \geq 1$,
there is a **set** \mathcal{U} of $k \cdot n^{\frac{1}{k}}$ **ultrametrics** and a mapping
home : $X \rightarrow \mathcal{U}$, such that for every $x, y \in U$,

$$d_{\text{home}(x)}(x, y) \leq (16 \cdot k) \cdot d_X(x, y)$$


Theorem (Ramsey based Deterministic Distance Oracle)

For any n -point metric space, there is a distance oracle with :

Distance Oracle	Distortion	Size	Query
This paper	$16 \cdot k$	$O(k \cdot n^{1+1/k})$	$O(1)$
This paper	$8(1 + \epsilon)k$	$O(n^{1+1/k})$	$O(1/\epsilon)$
This paper + C14	$2k - 1$	$O(k \cdot n^{1+1/k})$	$O(1)$
C15 (Randomized)	$2k - 1$	$O(n^{1+1/k})$	$O(1)$


Ramsey Spanning Tree Question

Given a **weighted graph** $G = (V, E, w)$, and a fixed distortion $k > 1$, what is the **largest subset** $M \subset V$, such that:
there is a **spanning tree** T of G with distortion k w.r.t $M \times V$?



Ramsey Spanning Tree Question

Given a **weighted graph** $G = (V, E, w)$, and a fixed distortion $k > 1$, what is the **largest subset** $M \subset V$, such that: there is a **spanning tree** T of G with distortion k w.r.t $M \times V$?



For all $v \in M$ and $u \in V$, $d_T(v, u) \leq k \cdot d_G(v, u)$.

Main Result

Ramsey Spanning Tree Question

Given a **weighted graph** $G = (V, E, w)$, and a fixed distortion $k > 1$, what is the **largest subset** $M \subset V$, such that:
there is a **spanning tree** T of G with distortion k w.r.t $M \times V$?

Theorem (Main Result)

*For every n -vertex **weighted graph** $G = (V, E, w)$ and $k \geq 1$,
there exists a **subset** M of size $n^{1-1/k}$ and **spanning tree** T of G
with **distortion** $O(k \cdot \log \log n)$ w.r.t $M \times V$.*

Main Result

Ramsey Spanning Tree Question

Given a **weighted graph** $G = (V, E, w)$, and a fixed distortion $k > 1$, what is the **largest subset** $M \subset V$, such that:
there is a **spanning tree** T of G with distortion k w.r.t $M \times V$?

Theorem (Main Result)

For every n -vertex **weighted graph** $G = (V, E, w)$ and $k \geq 1$,
there exists a subset M of size $n^{1-1/k}$ and **spanning tree** T of G
with distortion $O(k \cdot \log \log n)$ w.r.t $M \times V$.

Theorem (Mendel, Naor 07)

For every n -point **metric space** (X, d_X) and $k \geq 1$,
there exists a subset M of size $n^{1-1/k}$ and an **ultrametric** U over X
with distortion $O(k)$ w.r.t $M \times X$.

Theorem (Main Result)

For every n -vertex **weighted graph** $G = (V, E, w)$ and $k \geq 1$,
there exists a **subset** M of size $n^{1-1/k}$ and **spanning tree** T of G
with **distortion** $O(k \cdot \log \log n)$ w.r.t $M \times V$.


Theorem (Main Result)

For every n -vertex **weighted graph** $G = (V, E, w)$ and $k \geq 1$, there exists a **subset** M of size $n^{1-1/k}$ and **spanning tree** T of G with **distortion** $O(k \cdot \log \log n)$ w.r.t $M \times V$.

Corollary

For every n -vertex weighted graph $G = (V, E, w)$ and $k \geq 1$, there is a **set** \mathcal{T} of $k \cdot n^{\frac{1}{k}}$ **spanning trees** and a mapping $\text{home} : V \rightarrow \mathcal{T}$, such that for every $u, v \in V$,


$$d_{\text{home}(v)}(v, u) \leq O(k \cdot \log \log n) \cdot d_G(v, u)$$



Corollary


For every n -vertex weighted graph $G = (V, E, w)$ and $k \geq 1$, there is a **set \mathcal{T}** of $k \cdot n^{\frac{1}{k}}$ **spanning trees** and a mapping **home** : $V \rightarrow \mathcal{T}$, such that for every $u, v \in V$,

$$d_{\text{home}(v)}(v, u) \leq O(k \cdot \log \log n) \cdot d_G(v, u)$$




The union of all the trees in \mathcal{T} creates an $O(k \cdot \log \log n)$ -**spanner** with $O(k \cdot n^{1+\frac{1}{k}})$ edges.

Application: Compact Routing Scheme




- **Huge** network

Application: Compact Routing Scheme




- **Huge** network
- There is a **server** in each **node**.

Application: Compact Routing Scheme




- **Huge** network
- There is a **server** in each **node**.
- Task: **route** packages throughout the network.

Application: Compact Routing Scheme




- **Huge** network
- There is a **server** in each **node**.
- Task: **route** packages throughout the network.
- **Store** the whole network in each node is **unfeasible**.

Compact Routing Scheme




Initiating
node


Compact Routing Scheme




Compact Routing Scheme



Compact Routing Scheme




Compact Routing Scheme




In order to keep other **parameters small**, we will allow **stretch**.

Compact Routing Scheme



In order to keep other **parameters small**, we will allow **stretch**.
Stretch k : the length of a **route** from v to u will be $\leq k \cdot d_G(v, u)$.

Compact Routing Scheme



In order to keep other **parameters small**, we will allow **stretch**.
Stretch k : the length of a **route** from v to u will be $\leq k \cdot d_G(v, u)$.

Theorem (Thorup, Zwick, 01)


For any n -vertex **tree** $T = (V, E)$, there is a routing scheme with :

Stretch	Label	Table	Decision time
1	$O(\log n)$	$O(1)$	$O(1)$

Routing using Ramsey Spanning Trees

For every n -vertex weighted graph $G = (V, E, w)$ and $k \geq 1$, there is a **set** \mathcal{T} of $k \cdot n^{\frac{1}{k}}$ **spanning trees** and a mapping **home** : $V \rightarrow \mathcal{T}$, such that for every $u, v \in V$,

$$d_{\text{home}(v)}(v, u) \leq O(k \cdot \log \log n) \cdot d_G(v, u)$$




Routing using Ramsey Spanning Trees

For every n -vertex weighted graph $G = (V, E, w)$ and $k \geq 1$, there is a **set** \mathcal{T} of $k \cdot n^{\frac{1}{k}}$ **spanning trees** and a mapping **home** : $V \rightarrow \mathcal{T}$, such that for every $u, v \in V$,

$$d_{\text{home}(v)}(v, u) \leq O(k \cdot \log \log n) \cdot d_G(v, u)$$

To **route** a package from u to v , we will simply route on **home**(v)!




Routing using Ramsey Spanning Trees

For every n -vertex weighted graph $G = (V, E, w)$ and $k \geq 1$, there is a **set** \mathcal{T} of $k \cdot n^{\frac{1}{k}}$ **spanning trees** and a mapping **home** : $V \rightarrow \mathcal{T}$, such that for every $u, v \in V$,

$$d_{\text{home}(v)}(v, u) \leq O(k \cdot \log \log n) \cdot d_G(v, u)$$

To **route** a package from u to v , we will simply route on **home**(v)!

The label of v will consist of: $(\text{home}(v), \text{Label}_{\text{home}(v)}(v))$.



Routing using Ramsey Spanning Trees


For every n -vertex weighted graph $G = (V, E, w)$ and $k \geq 1$, there is a **set** \mathcal{T} of $k \cdot n^{\frac{1}{k}}$ **spanning trees** and a mapping **home** : $V \rightarrow \mathcal{T}$, such that for every $u, v \in V$,

$$d_{\text{home}(v)}(v, u) \leq O(k \cdot \log \log n) \cdot d_G(v, u)$$

To **route** a package from u to v , we will simply route on **home**(v)!

The label of v will consist of: $(\text{home}(v), \text{Label}_{\text{home}(v)}(v))$.

The table of v will consist of union of all tables in \mathcal{T} .



Routing using Ramsey Spanning Trees

For every n -vertex weighted graph $G = (V, E, w)$ and $k \geq 1$, there is a **set** \mathcal{T} of $k \cdot n^{\frac{1}{k}}$ **spanning trees** and a mapping **home** : $V \rightarrow \mathcal{T}$, such that for every $u, v \in V$,

$$d_{\text{home}(v)}(v, u) \leq O(k \cdot \log \log n) \cdot d_G(v, u)$$

To **route** a package from u to v , we will simply route on **home**(v)!

The label of v will consist of: $(\text{home}(v), \text{Label}_{\text{home}(v)}(v))$.

The table of v will consist of union of all tables in \mathcal{T} .

Theorem (Ramsey based Compact Routing Scheme)

For any n -vertex graph, there is a routing scheme with :

Stretch	Label	Table	Decision time
$O(k \cdot \log \log n)$	$O(\log n)$	$O(k \cdot n^{\frac{1}{k}})$	$O(1)$

Theorem (Ramsey based Compact Routing Scheme)

For any n -vertex graph, there is a routing scheme with :

Stretch	Label	Table	Decision time
$O(k \cdot \log \log n)$	$O(\log n)$	$O(k \cdot n^{\frac{1}{k}})$	$O(1)$

Theorem (Thorup, Zwick 01, Chechik 13)

For any n -vertex graph, there is a routing scheme with :

Stretch	Label	Table	Decision time
$3.68k = O(k)$	$O(k \cdot \log n)$	$O(k \cdot n^{\frac{1}{k}})$	$O(1)$ (initial: $O(k)$)

Theorem (Ramsey based Compact Routing Scheme)

For any n -vertex graph, there is a routing scheme with :

Stretch	Label	Table	Decision time
$O(k \cdot \log \log n)$	$O(\log n)$	$O(k \cdot n^{\frac{1}{k}})$	$O(1)$

Theorem (Thorup, Zwick 01, Chechik 13)

For any n -vertex graph, there is a routing scheme with :

Stretch	Label	Table	Decision time
$3.68k = O(k)$	$O(k \cdot \log n)$	$O(k \cdot n^{\frac{1}{k}})$	$O(1)$ (initial: $O(k)$)

By choosing $k = \log n$, we get:

	Stretch	Label	Table	D. time
Here	$O(\log n \cdot \log \log n)$	$O(\log n)$	$O(\log n)$	$O(1)$
[TZ01]	$O(\log n)$	$O(\log^2 n)$	$O(\log n)$	$O(1)$ ($O(\log n)$)

Technical Ideas

Theorem (Main Result)

For every n -vertex **weighted graph** $G = (V, E, w)$ and $k \geq 1$,
there exists a **subset** M of size $n^{1-1/k}$ and **spanning tree** T of G
with **distortion** $O(k \cdot \log \log n)$ w.r.t $M \times V$.

Technical Ideas

Theorem (Main Result)

For every n -vertex **weighted graph** $G = (V, E, w)$ and $k \geq 1$, there exists a **subset** M of size $n^{1-1/k}$ and **spanning tree** T of G with **distortion** $O(k \cdot \log \log n)$ w.r.t $M \times V$.

- Framework: Petal decomposition.

Technical Ideas

Theorem (Main Result)

For every n -vertex **weighted graph** $G = (V, E, w)$ and $k \geq 1$, there exists a **subset** M of size $n^{1-1/k}$ and **spanning tree** T of G with **distortion** $O(k \cdot \log \log n)$ w.r.t $M \times V$.

- Framework: Petal decomposition.
- Hierarchically padded decompositions.


Technical Ideas

Theorem (Main Result)


For every n -vertex **weighted graph** $G = (V, E, w)$ and $k \geq 1$, there exists a **subset** M of size $n^{1-1/k}$ and **spanning tree** T of G with **distortion** $O(k \cdot \log \log n)$ w.r.t $M \times V$.

- Framework: Petal decomposition.
- Hierarchically padded decompositions.
- Region growing.


Petal Decomposition




Petal Decomposition




Petal Decomposition




Petal Decomposition




Petal Decomposition




Petal Decomposition




Petal Decomposition




Petal Decomposition




Petal Decomposition



Petal Decomposition




Petal Decomposition




Petal Decomposition

- Each cluster X (petal) has a center vertex x .




Petal Decomposition

- Each cluster X (petal) has a center vertex x .
- The radius Δ defined w.r.t the center.




Petal Decomposition

- Each cluster X (petal) has a center vertex x .
- The radius Δ defined w.r.t the center.
- The radius decrease by $\frac{3}{4}$ factor in each hierarchi. step.




Petal Decomposition

- Each cluster X (petal) has a center vertex x .
- The radius Δ defined w.r.t the center.
- The radius decrease by $\frac{3}{4}$ factor in each hierarchi. step.
- The radius of T is at most 4 times larger than in G .



Petal Decomposition

- Each cluster X (petal) has a center vertex x .
- The radius Δ defined w.r.t the center.
- The radius decrease by $\frac{3}{4}$ factor in each hierarchi. step.
- The radius of T is at most 4 times larger than in G .




Corollary

Suppose v, u were separated while being in cluster of radius Δ . Then $d_T(v, u) \leq 8 \cdot \Delta$.

Petal Growth

Degree of freedom:

parameter $R \in [\text{lo}, \text{hi}]$ ($\text{hi} - \text{lo} = \frac{\Delta}{8}$).




Petal Growth

Degree of freedom:

parameter $R \in [\text{lo}, \text{hi}]$ ($\text{hi} - \text{lo} = \frac{\Delta}{8}$).

W_r denotes the **petal** (cluster)
created for $R = r$.

Monotonicity: $r' \leq r \Rightarrow W_{r'} \subseteq W_r$.



Petal Growth


Degree of freedom:

parameter $R \in [\text{lo}, \text{hi}]$ ($\text{hi} - \text{lo} = \frac{\Delta}{8}$).

W_r denotes the **petal** (cluster)
created for $R = r$.

Monotonicity: $r' \leq r \Rightarrow W_{r'} \subseteq W_r$.

Set $\delta = \Delta / (k \cdot \log \log n)$.



Petal Growth

Degree of freedom:


parameter $R \in [\text{lo}, \text{hi}]$ ($\text{hi} - \text{lo} = \frac{\Delta}{8}$).

W_r denotes the **petal** (cluster)
created for $R = r$.

Monotonicity: $r' \leq r \Rightarrow W_{r'} \subseteq W_r$.

Set $\delta = \Delta / (k \cdot \log \log n)$.

Vertex v s.t. $B(v, \delta) \subseteq W_r$ is **padding**.



Petal Growth

Degree of freedom:

parameter $R \in [\text{lo}, \text{hi}]$ ($\text{hi} - \text{lo} = \frac{\Delta}{8}$).


W_r denotes the **petal** (cluster)
created for $R = r$.

Monotonicity: $r' \leq r \Rightarrow W_{r'} \subseteq W_r$.

Set $\delta = \Delta / (k \cdot \log \log n)$.

Vertex v s.t. $B(v, \delta) \subseteq W_r$ is **padded**.

All vertices out of $W_{r+\delta} \setminus W_{r-\delta}$
(restricted area) are padded.



Petal Growth

Degree of freedom:

parameter $R \in [\text{lo}, \text{hi}]$ ($\text{hi} - \text{lo} = \frac{\Delta}{8}$).


W_r denotes the **petal** (cluster) created for $R = r$.

Monotonicity: $r' \leq r \Rightarrow W_{r'} \subseteq W_r$.

Set $\delta = \Delta / (k \cdot \log \log n)$.

Vertex v s.t. $B(v, \delta) \subseteq W_r$ is **padded**.

All vertices out of $W_{r+\delta} \setminus W_{r-\delta}$
(restricted area) are padded.



Padded vertices suffer distortion **at most** $\Delta/\delta = O(k \cdot \log \log n)!$

Petal Growth

Degree of freedom:

parameter $R \in [\text{lo}, \text{hi}]$ ($\text{hi} - \text{lo} = \frac{\Delta}{8}$).


W_r denotes the **petal** (cluster)
created for $R = r$.

Monotonicity: $r' \leq r \Rightarrow W_{r'} \subseteq W_r$.

Set $\delta = \Delta / (k \cdot \log \log n)$.

Vertex v s.t. $B(v, \delta) \subseteq W_r$ is **padded**.


All vertices out of $W_{r+\delta} \setminus W_{r-\delta}$
(**restricted area**) are padded.



Padded vertices suffer distortion **at most** $\Delta/\delta = O(k \cdot \log \log n)!$


Goal: find r , with **many padded vertices!** (**sparse restricted area**).

Petal Decomposition



Petal Decomposition


A vertex which is **padded in all the levels** will have small distortion w.r.t all other vertices.



Petal Decomposition

A vertex which is **padded in all the levels** will have small distortion w.r.t all other vertices.

Goal: choose parameters $(r \in [lo, hi])$ s.t. at least $n^{1-\frac{1}{k}}$ vertices will be **padded in all levels**.




Petal Decomposition

A vertex which is **padded in all the levels** will have small distortion w.r.t all other vertices.

Goal: choose parameters $(r \in [lo, hi])$ s.t. at least $n^{1-\frac{1}{k}}$ vertices will be **padded in all levels**.

A vertex is called **active** if it is **padded in all levels up till now**.




Region Growing

For petal W_r :

Active $x \in W_{r-\delta}$ **remains** active.

Active $x \in W_{r+\delta} \setminus W_{r-\delta}$
ceases to be active.




Region Growing

For petal W_r :

Active $x \in W_{r-\delta}$ **remains** active.

Active $x \in W_{r+\delta} \setminus W_{r-\delta}$
ceases to be active.




Region Growing

For petal W_r :

Active $x \in W_{r-\delta}$ **remains** active.

Active $x \in W_{r+\delta} \setminus W_{r-\delta}$
ceases to be active.




Region Growing

For petal W_r :

Active $x \in W_{r-\delta}$ **remains** active.

Active $x \in W_{r+\delta} \setminus W_{r-\delta}$
ceases to be active.




Region Growing

For petal W_r :

Active $x \in W_{r-\delta}$ **remains** active.

Active $x \in W_{r+\delta} \setminus W_{r-\delta}$
ceases to be active.



Intuition

*There is $r \in [lo, hi]$ such that $W_{r-\delta}$ is
large enough compared to $W_{r+\delta}$.*

Intuition

There is $r \in [lo, hi]$ such that $W_{r-\delta}$ is
large enough *compared to $W_{r+\delta}$.*

Intuition

There is $r \in [lo, hi]$ such that $W_{r-\delta}$ is
large enough compared to $W_{r+\delta}$.



Intuition

There is $r \in [lo, hi]$ such that $W_{r-\delta}$ is
large enough compared to $W_{r+\delta}$.

Corollary

At least $n^{1-1/k}$ vertices **remain active** at the **end** of the process.




Intuition

There is $r \in [lo, hi]$ such that $W_{r-\delta}$ is
large enough compared to $W_{r+\delta}$.

Corollary

At least $n^{1-1/k}$ vertices **remain active** at the **end** of the process.



Theorem (Main Result)

For every n -vertex **weighted graph** $G = (V, E, w)$ and $k \geq 1$,
there exists a **subset** M of size $n^{1-1/k}$ and **spanning tree** T of G
with **distortion** $O(k \cdot \log \log n)$ w.r.t $M \times V$.

Open Questions

- 1 **Remove** the $\log \log n$ factor.

Conjecture

For every n -vertex **weighted graph** $G = (V, E, w)$ and $k \geq 1$, there exists a **subset** M of size $n^{1-1/k}$ and **spanning tree** T of G with **distortion** $O(k \cdot \log \log n)$ w.r.t $M \times V$.

Open Questions

- 1 **Remove** the $\log \log n$ factor.

Conjecture

For every n -vertex **weighted graph** $G = (V, E, w)$ and $k \geq 1$, there exists a **subset** M of size $n^{1-1/k}$ and **spanning tree** T of G with **distortion** $O(k \cdot \log \log n)$ w.r.t $M \times V$.

- 2 Improve construction for deterministic distance oracle.

Distance Oracle	Distortion	Size	Query
This paper +C14	$2k - 1$	$O(k \cdot n^{1+1/k})$	$O(1)$
C15 (Randomized)	$2k - 1$	$O(n^{1+1/k})$	$O(1)$

Open Questions

- 1 **Remove** the $\log \log n$ factor.

Conjecture

For every n -vertex **weighted graph** $G = (V, E, w)$ and $k \geq 1$, there exists a **subset** M of size $n^{1-1/k}$ and **spanning tree** T of G with **distortion** $O(k \cdot \log \log n)$ w.r.t $M \times V$.

- 2 Improve construction for deterministic distance oracle.

Distance Oracle	Distortion	Size	Query
This paper +C14	$2k - 1$	$O(k \cdot n^{1+1/k})$	$O(1)$
C15 (Randomized)	$2k - 1$	$O(n^{1+1/k})$	$O(1)$

- 3 Find more **applications** to Ramsey spanning trees!